arXiv:1401.7462v3 [math.GR] 29 Sep 2014

On element orders in covers of finite simple groups of Lie type

Mariya A. Grechkoseeva[1]

1. Introduction

Given a finite group G, we say that a finite group H is a *cover* of G if G is a homomorphic image of H. Denoting by $\omega(A)$ the set of element orders of a group A and referring to it as the *spectrum* of A, we say that G is *recognizable (by spectrum) among covers* if $\omega(G) \neq \omega(H)$ for any proper cover H of G. If G has a nontrivial normal soluble subgroup, then G is not recognizable among covers, and in fact, there are infinitely many nonisomorphic covers H of G with $\omega(H) = \omega(G)$ [1, Lemma 1].

The present paper concerns recognizability of finite nonabelian simple groups, and we denote these groups according to the 'Atlas of finite groups' [2]. The following simple groups are known to be recognizable by spectrum among covers: the sporadic groups (see [3]), the alternating groups [4], the Ree and Suzuki groups [5–7], the exceptional groups of types G_2 and E_8 [8, 9], the groups $L_2(q)$ [4, 10], the groups $L_3(q)$ [11, 12], the groups $L_n(q)$ with $n \geqslant 5$ [13], and the groups $U_n(q)$, with $n \geqslant 5$, other than $U_5(2)$ [14]. Also it is worth noting that $L_4(q)$ and $U_4(q)$ are recognizable among covers as long as q is even or prime [13–15], and $U_3(q)$ is not recognizable among covers if and only if q is a Mersenne prime with $q^2 - q + 1$ being a prime too [16]. Lastly, $L_4(13^{24})$, $U_5(2)$ and ${}^3D_4(2)$ are not recognizable among covers [14, 17, 18].

Our purpose is not to give an exhaustive answer to the question what finite simple groups of Lie type are recognizable by spectrum among covers and which are not but rather to show that groups of sufficiently large Lie rank have this property of recognizability. Thus we do not touch the very interesting case of the groups $L_4(q)$ and $U_4(q)$ but focus our attention on groups not mentioned above. Recall that $O_{2n+1}(q) \simeq S_{2n}(q)$ provided that $n = 2$ or q is even.

Theorem 1.1. *If S is one of the simple groups ${}^3D_4(q)$ with $q > 2$, $F_4(q)$, $E_6(q)$, ${}^2E_6(q)$, $E_7(q)$, $S_{2n}(q)$ with $n \geqslant 2$, $O_{2n+1}(q)$ with $n \geqslant 3$ and q odd, and $O^{\pm}_{2n}(q)$ with $n \geqslant 4$, then S is recognizable by spectrum among covers.*

The theorem and before-mentioned results yield the following corollary.

Corollary 1.1. *If S is a finite nonabelian simple group other than $L_4(q)$, $U_3(q)$, $U_4(q)$, $U_5(2)$, and ${}^3D_4(2)$, then S is recognizable by spectrum among covers.*

Our strategy of proof is standard and based on the following two observations (in these observations, as well as throughout the paper, all modules and representations are finite-dimensional). First, it is not hard to see that a group G is recognizable by spectrum among covers as long as $\omega(G) \neq \omega(V \leftthreetimes G)$ for any nontrivial G-module V over a field of positive characteristic (cf. Lemma 2.3 below). Second, given a finite group of Lie type, it is natural to distinguish representations in the defining characteristic from other representations. Thus for every simple group S satisfying the hypothesis of the theorem and every nontrivial S-module V, we need to prove that $\omega(S) \neq \omega(V \leftthreetimes S)$ provided that the corresponding representation (i) is equicharacteristic, and (ii) is cross-characteristic (cf. Problems 17.73 and 17.74 in [19]).

[1]Supported by the RFBR (12-01-31221 and 12-01-90006).

Proposition 1.1. *Let q be a power of a prime p and let S be one of the simple groups ${}^3D_4(q)$ with $q > 2$, $F_4(q)$, $E_6(q)$, ${}^2E_6(q)$, $E_7(q)$, $S_{2n}(q)$ with $n \geqslant 2$, $O_{2n+1}(q)$ with $n \geqslant 3$ and q odd, and $O^{\pm}_{2n}(q)$ with $n \geqslant 4$. If V is a nonzero S-module over a field of characteristic p and H is a natural semidirect product of V and S, then $\omega(S) \neq \omega(H)$.*

Under condition (ii), the problem has already been settled for symplectic and orthogonal groups of dimension at least six [14]. So we deal with the remaining groups.

Proposition 1.2. *Let S be one of the simple groups ${}^3D_4(q)$, $F_4(q)$, $E_6(q)$, ${}^2E_6(q)$, $E_7(q)$, and $S_4(q)$. If V is a nonzero S-module over a field of characteristic r, where r is coprime to q, and H is a natural semidirect product of V and S, then $\omega(S) \neq \omega(H)$.*

Proposition 1.1 and [8] give an affirmative answer to [19, Problem 17.73] in all cases except for the cases (a) and (f). In the case (f) the answer is negative [18] and the case (a) concerns $U_4(q)$. Proposition 1.2, [14, Theorem 1] and [13, Lemma 12] give an affirmative answer to [19, Problem 17.74].

We conclude this paper with a result comparing the spectrum of a proper cover of $O_{2n+1}(q)$ with the spectrum of $S_{2n}(q)$. This result is motivated by the following assertion: if $L = S_{2n}(q)$, where $n \geqslant 4$, and G is a finite group such that $\omega(G) = \omega(L)$, then G has a unique nonabelian composition factor and if this factor is a group of Lie type in the same characteristic as L, then it is isomorphic either to L or to one of the groups $O_{2n+1}(q)$ and $O^-_{2n}(q)$ [20, Theorem 3]. It is well-known that the spectra of $S_{2n}(q)$ and $O_{2n+1}(q)$ are very close. This property of $S_{2n}(q)$ and $O_{2n+1}(q)$ makes the case when the factor is isomorphic to $O_{2n+1}(q)$ more difficult to eliminate. Somewhat paradoxically, the same property allows us to apply our results on covers to this situation.

Proposition 1.3. *Let $L = S_{2n}(q)$ and $S = O_{2n+1}(q)$, where $n \geqslant 3$ and q is odd. If H is a proper cover of S, then $\omega(H) \not\subseteq \omega(L)$. In particular, if G is a finite group such that $\omega(G) = \omega(L)$ and G has a composition factor isomorphic to S, then $S \leqslant G \leqslant \mathrm{Aut}(S)$.*

2. Preliminaries

Our notation for classical groups and groups of Lie type is based on that in [21] with the following exception: $\Phi(q)$ always stands for the adjoint version, while the universal version is denoted by $\Phi(q)_u$. The same rule applies to the symbol ${}^d\Phi(q)$. Also sometimes we substitute $\Phi(q)$ and ${}^2\Phi(q)$ by $\Phi^+(q)$ and $\Phi^-(q)$ respectively .

Lemma 2.1 (Zsigmondy [22])**.** *Let $q \geqslant 2$ and $n \geqslant 3$ be integers, with $(q, n) \neq (2, 6)$. There exists a prime r such that r divides $q^n - 1$ but does not divide $q^i - 1$ for $i < n$.*

With notation of Lemma 2.1, we call a prime r a *primitive prime divisor* of $q^n - 1$ and denote it by $r_n(q)$.

By (n, m) we denote the greatest common divisor of two positive integers n and m. Given a positive integer n, we write $\pi(n)$ for the set of all prime divisors of n. Also, if r is another positive integer, then n_r denotes the $\pi(r)$-part of n (that is, the largest divisor m of n such that $\pi(m) \subseteq \pi(r)$), and $n_{r'}$ denotes the ratio n/n_r.

If G is a finite group and r is a prime, we refer to the highest power of r that lies in $\omega(G)$ as the r-*exponent* of G. If G is a group of Lie type over a field of positive characteristic p, then it is natural and instructive to consider $\omega(G)$ as a disjoint union of the following three subsets:

the subset of numbers that are powers of p, the subset $\omega_{p'}(G)$ of numbers that are coprime to p, and the subset $\omega_{mix}(G)$ of the rest, so called 'mixed', orders.

If G is a finite group, then $\pi(G) = \pi(|G|)$. The *prime graph* of G is a graph whose vertex set is $\pi(G)$ and in which two different vertices are adjacent if and only if their product lies in $\omega(G)$. The following property of the simple groups of Lie type is well-known but rarely appears as a separate assertion, so we include the corresponding lemma in this paper.

Lemma 2.2. *Suppose that S is a finite simple group of Lie type. Then for every $r \in \pi(S)$, there is $t \in \pi(S)$ such that $r \neq t$ and $rt \notin \omega(S)$.*

Proof. If S is one of the simple groups $L_2(q)$, $L_3(q)$, $U_3(q)$, and $S_4(q)$, or S is a simple exceptional group of type other than E_7, then by [23,24], the prime graph of S is disconnected, and the assertion follows. For the rest of simple classical groups, the lemma is proved in [14, Lemma 12]. If $S = E_7(q)$, then we can take t to be equal to $r_9(q)$ or $r_{18}(q)$ (see, for example, [25, Fig. 4]).

The next lemma is also well-known and in the special case $A = B$, is proved, for example, in [11, Lemma 12].

Lemma 2.3. *Let A and B be finite groups. The following are equivalent.*

(1) *$\omega(H) \not\subseteq \omega(B)$ for any proper cover H of A;*

(2) *$\omega(H) \not\subseteq \omega(B)$ for any split extension $H = K : A$, where K is a nontrivial elementary abelian group.*

Proof. Clearly, (1) implies (2). Suppose that (2) does not imply (1), and let H be a group of minimum possible order such that H is a proper cover of A and $\omega(H) \subseteq \omega(B)$. Let K be a normal subgroup of H with $H/K = A$. Choose $r \in \pi(K)$ and let R be a Sylow r-subgroup of K. If $N = N_H(R)$, then the Frattini argument yields $A \simeq H/K = NK/K \simeq N/N \cap K$. Since $N \cap K \neq 1$ and $\omega(N) \subseteq \omega(H)$, we have $H = N$ by the minimality of H. Repeating this observation for other prime divisors of $|K|$, we conclude that K is nilpotent. Denote the Hall r'-subgroup of K by C and the Frattini subgroup of R by $\Phi(R)$. The factor group of $H/(C \times \Phi(R))$ by $K/(C \times \Phi(R))$ is isomorphic to A, and again the minimality of H yields $C = \Phi(R) = 1$. Thus K is an elementary abelian r-group, and in particular, the group $A = H/K$ acts on K by conjugation. If H_1 is a semidirect product of K and A with respect to this action, then $\omega(H_1) \subseteq \omega(H) \subseteq \omega(B)$ by [4, Lemma 10]. This contradicts (2).

Lemma 2.4 ([26, Lemma 1])**.** *Let G be a finite group, K a normal subgroup of G and let G/K be a Frobenius group with kernel F and cyclic complement C. If $(|F|, |K|) = 1$ and F is not contained in $KC_G(K)/K$, then $r|C| \in \omega(G)$ for some prime divisor r of $|K|$.*

Lemma 2.5. *The group $SL_n(q)$, where $n \geqslant 2$, includes a Frobenius subgroup with kernel of order q^{n-1} and cyclic complement of order $(q^{n-1}-1)_{(n,q-1)'}$, as well as a Frobenius subgroup with kernel of order q^k and cyclic complement of order $q^k - 1$ for every $1 \leqslant k < n-1$.*

Proof. It is well known that the natural semidirect product of a vector space of order q^k by the Zinger cycle of order $q^k - 1$ in $GL_k(q)$ is a Frobenius group. It is easily seen that this product can be embedded into a line stabilizer in $GL_{k+1}(q)$. Since $GL_{k+1}(q)$, where $1 \leqslant k < n-1$, can be embedded into $SL_n(q)$, the second assertion follows. Furthermore, $PSL_n(q)$ includes a Frobenius subgroup F with kernel of order q^{n-1} and cyclic complement of order $(q^{n-1}-1)/d$,

where $d = (n, q-1)$ (see, for example, [27, Lemma 5]). Denoting $\pi = \pi(d)$ we see that the full preimage of the Hall π'-subgroup of F in $SL_n(q)$ is a split extension of a group of order d by the desired Frobenius group of order $q^{n-1}(q^{n-1}-1)_{d'}$.

Lemma 2.6. *The simple group $S_{2n}(q)$, where q is even and $n = 2^m \geqslant 2$, includes a Frobenius subgroup with cyclic kernel of order $q^n + 1$ and cyclic complement of order $2n$.*

Proof. The proof is similar to that of Lemma 2.4 in [28].

Lemma 2.7. *Let S be a finite simple group of Lie type over a field of characteristic p and let S acts faithfully on a vector space V over a field of characteristic r, where $r \neq p$. Let $H = V \leftthreetimes S$ be a natural semidirect product of V by S. Suppose that s is a power of r and that s lies in the spectrum of some proper parabolic subgroup P of S. If the unipotent radical of P is abelian or both p and r are odd or $p = 2$ and r is not a Fermat prime or $r = 2$ and p is not a Mersenne prime, then $rs \in \omega(H)$.*

Proof. Let $g \in P$ be an element of order s, U a unipotent radical of P and let $M = U \leftthreetimes \langle g \rangle$. By [21, Theorem 2.6.5e] it follows that $C_G(U) = Z(U)$. Thus $O_r(M) = 1$, and by the Hall–Higman theorem [29, Theorem 2.1.1], the degree of the minimal polynomial of g on V is equal to s. This implies that the coset Vg contains an element of order rs.

3. SR-subgroups of finite groups of Lie type

Let G be a finite universal group of Lie type. By [21, Theorem 2.2.6e], there are a simply connected simple algebraic group $\overline{G}$ and a Frobenius endomorphism σ of $\overline{G}$ such that $G = C_{\overline{G}}(\sigma)$. By [21, Theorem 2.1.6], there is a σ-stable maximal torus $\overline{T}$ of $\overline{G}$. Let Φ denote the root system of $\overline{G}$ with respect to $\overline{T}$ and Π denote some fundamental system. Also we write $\overline{X}_\alpha$ to denote a root subgroup of $\overline{G}$ associated with a root α. Let $\Delta \subseteq \Pi$. Following [30], we call a subgroup of $\overline{G}$ an *SR-subgroup* (with respect to $\overline{T}$) if it is generated by the subgroups $\overline{X}_{\pm\alpha}$ for $\alpha \in \Delta$. A subgroup of the finite group G is an SR-subgroup if it is an intersection of an SR-subgroup of $\overline{G}$ with G.

Lemma 3.1. *Let G be a finite universal group of Lie type over a field of characteristic p and let $g \in G$ lie in some SR-subgroup of type Ψ. The element g has an nonzero fixed vector in every nonzero G-module over a field of characteristic p in the following cases:*

(a) $G = A_n(q)_u$, $n \geqslant 1$, *and* $\Psi = A_m$, *where* $m \leqslant n/2$;

(b) $G = B_n(q)_u$, $n \geqslant 3$, q *is odd, and* $\Psi = A_{n-1}$;

(c) $G = C_n(q)_u$, $n \geqslant 2$, q *is odd, and* $\Psi = C_{n-1}$;

(d) $G = D_n^{\pm}(q)_u$, $n \geqslant 4$, *and* $\Psi = A_{n-2}$;

(e) $G = E_7(q)_u$ *and* $\Psi = E_6$;

(f) $G = E_6^{\pm}(q)_u$ *and* $\Psi = D_5$.

Proof. See Corollaries 1.6, 1.7, 4.3, 5.1, and also a remark after Corollary 1.3 in [30].

Table 1: Generators of $Z(\overline{G})$ for some $\overline{G}$

Φ	Dynkin diagram	generators of $Z(\overline{G})$
B_n		$h_n(-1)$
C_n		$h_1(-1)h_3(-1)\dots h_n((-1)^n)$
D_n, n even		$h_1(-1)h_3(-1)\dots h_{n-1}(-1)$, $h_{n-1}(-1)h_n(-1)$
D_n, n odd		$h_1(-1)h_3(-1)\dots h_{n-2}(-1)h_{n-1}(t)h_n(t)$, $t^2=-1$
E_6		$h_1(t)h_2(t^2)h_5(t)h_6(t^2)$, $t^3=1$
E_7		$h_4(-1)h_5(-1)h_7(-1)$

We need an easy consequence of Lemma 3.1 for simple groups. For convenience, we list generators of the centers of some simple algebraic groups from [21, Table 1.12.6]. In Table 1, the symbol $\overline{G}$ denotes a simple algebraic group of type Φ, and $h_i(t)$ denotes $h_{\alpha_i}(t)$.

Lemma 3.2. *Let S be a finite simple groups of Lie type over a field of characteristic p, V a nonzero S-module over a field of the same characteristic p and H the natural semidirect product of V by S. Then $\omega(H)$ includes the set $p\cdot\omega_{p'}(\Sigma)$, where*

(a) $\Sigma = SL_n(q)$ *if* $S = O_{2n+1}(q)$, $n \geqslant 3$, q *is odd or* $S = O^{\pm}_{2n+2}(q)$, $n \geqslant 3$;

(b) $\Sigma = SU_4(q)$ *if* $S = O_9(q), O^{\pm}_{10}(q)$, q *is odd;*

(c) $\Sigma = E_6(q)_u$ *if* $S = E_7(q)$;

(d) $\Sigma = Spin_{10}(q)$ *if* $S = E_6(q)$;

(e) $\Sigma = Spin^{-}_{8}(q)$ *if* $S = {}^2E_6(q)$.

Proof. Let G be a finite universal group of Lie type such that $G/Z(G) = S$. We may assume that V is a G-module. We prove the lemma by finding an SR-subgroup Γ of G such that $\Gamma \simeq \Sigma$, $\Gamma \cap Z(G) = 1$ and every element of Γ has a nonzero fixed vector in V.

Since SR-subgroups are a special case of subsystem subgroups, we need some information on the subsystem subgroups from [21, Section 2.6]. Let $\overline{G}$, σ, $\overline{T}$, Φ, and Δ have the same meaning as those in the beginning of this section. Observe that $Z(G) = Z(\overline{G}) \cap G$ by [21, Proposition 2.5.9]. Let τ be the isometry of order 2 of the system E_6 induced by the symmetry of the Dynkin diagram if $S = {}^2E_6(q)$, and take $\tau = 1$ in other cases. Supposing that Δ generates closed, irreducible and τ-invariant subsystem Ψ of Φ, we define $\overline{\Gamma}_\Delta = \langle \overline{X}_\alpha \mid \alpha \in \Psi\rangle$. Since $\langle \overline{X}_\alpha \mid \alpha \in \Psi\rangle = \langle \overline{X}_\alpha, \overline{X}_{-\alpha} \mid \alpha \in \Delta\rangle$ by [21, Theorem 1.12.7d], it follows that $\overline{\Gamma}_\Delta$ is an SR-subgroup. Let W and W_0 be the Weyl groups of Φ and Ψ respectively, and let W_0^* be the stabilizer in W of Ψ. For every $w \in W_0^*$ we define an SR-subgroup $\Gamma_{\Delta,w}$ as follows. Let $\overline{N} = N_{\overline{G}}(\overline{T})$ and choose $n \in \overline{N}$ mapping to w. Then choose $x \in \overline{G}$ so that $\sigma = (\sigma n)^x$ and set

$$\Gamma_{\Delta,w} = C_{\overline{\Gamma}^x_\Delta}(\sigma).$$

If τw induces an isometry of Ψ of order d_1, then $\Gamma_{\Delta,w}$ is the universal group of type ${}^{d_1}\Psi(q)$ by [21, Proposition 2.6.2b,d].

Let $S = O_{2n+1}(q)$, where $n \geqslant 3$ and q is odd. Take $\Delta = \{\alpha_1, \alpha_2, \ldots, \alpha_{n-1}\}$ with notation for roots as in Table 1 and set $\Gamma = \Gamma_{\Delta,1}$. Then every element of Γ has a nonzero fixed vector in V by Lemma 3.1. Also it follows by the previous paragraph that $\Gamma = A_{n-1}(q)_u = SL_n(q)$. Lastly, $Z(G) = \langle h_{\alpha_n}(-1)\rangle$ and G is universal, therefore, $\Gamma \cap Z(G) = 1$. The groups $O^{\pm}_{2n+2}(q)$, $n \geqslant 3$, $E_7(q)$, $E_6(q)$ and ${}^2E_6(q)$ can be handled similarly with subsystems of types A_{n-1}, E_6, D_5 and D_4, respectively, in place of Ψ.

Let S be as in (b). Since $O_9(q)$, where q is odd, can be embedded into $O^{\pm}_{10}(q)$, it suffices to consider the case $S = O_9(q)$. Let $\Delta = \{\alpha_1, \alpha_2, \alpha_3\}$ (again with notation of Table 1). It is not hard to see that $|W_0^*/W_0| = 2$ and some $w \in W_0^* \setminus W_0$ induces an isometry of Ψ of order two. Taking $\Gamma = \Gamma_{\Delta,w}$ and reasoning as before, we conclude that $\Gamma = {}^2A_3(q)_u = SU_4(q)$, every element of Γ has a nonzero fixed vector in V, and $\Gamma \cap Z(G) = 1$. This completes the proof.

4. Proof of Theorem 1.1

We begin work by proving Proposition 1.1. Following [31], we call a finite group L of Lie type over a field of characteristic p *unisingular* if for every nontrivial abelian p-group A with L-action, every element of L has a nontrivial fixed point on A.

Proposition 1.1. *Let q be a power of a prime p and let S be one of the simple groups ${}^3D_4(q)$ with $q > 2$, $F_4(q)$, $E_6(q)$, ${}^2E_6(q)$, $E_7(q)$, $S_{2n}(q)$ with $n \geqslant 2$, $O_{2n+1}(q)$ with $n \geqslant 3$ and q odd, and $O^{\pm}_{2n}(q)$ with $n \geqslant 4$. If V is a nonzero S-module over a field of characteristic p and H is a natural semidirect product of V and S, then $\omega(S) \neq \omega(H)$.*

Proof. If the action of S on V is not faithful, then $C_S(V)$ is a nontrivial normal subgroup of S, and hence $C_S(V) = S$. This implies that for every prime $t \in \pi(S) \setminus \{p\}$, the product pt lies in $\omega(H)$, and so $\omega(H) \neq \omega(S)$ by Lemma 2.2. Thus we can assume that S acts faithfully on V.

Let $S = F_4(q)$. By [32], we know that $q^4 - q^2 + 1 \in \omega(S)$ and $p(q^4 - q^2 + 1) \notin \omega(S)$. Since S is unisingular [33, Theorem 1.3], it follows that $p(q^4 - q^2 + 1) \in \omega(H) \setminus \omega(S)$.

Let $S = {}^3D_4(q)$, where $q > 2$. It is well known that S includes a subgroup isomorphic to $G_2(q)$ (see, for example, [34, Table 8.51]). If q is odd, then $G_2(q)$ is unisingular [33, Theorem 1.3], and again using [32], we conclude that $p(q^2 - 1) \in \omega(H) \setminus \omega(S)$. Now let q be even. We work similarly to the proof of Lemma 3.1 in [8]. This proof involves three subgroups A, B and C of $G_2(q)$ such that $B \simeq SL_3(q)$, A is an SR-subgroup of type A_1 in B, $C \simeq SL_2(q)$ and C centralizes A. We choose $g \in A$ to be an element of order $q + 1$. Since g has a nonzero fixed vector in any B-module over a field of characteristic 2 by Lemma 3.1, it follows that $W = C_V(g) \neq 0$. The group C acts on W and includes a Frobenius subgroup with cyclic kernel F of order $q - 1$ and complement of order 2. If $F \subseteq C_C(W)$, then H has an element of order $2(q^2 - 1)$. Otherwise, we apply Lemma 2.4 and derive that H has an element of order $4(q+1)$. The centralizers of semisimple elements of S are described in [35], and their structure implies that $\omega_{mix}(S)$ consists of the divisors of the numbers $2(q^3 + 1)$, $2(q^3 - 1)$, $4(q^2 + q + 1)$, and $4(q^2 - q + 1)$. Since $q > 2$, it follows that neither $2(q^2 - 1)$ nor $4(q + 1)$ divides any of these numbers, and hence $\omega(H) \neq \omega(S)$.

Let $S = E^{\pm}_6(q)$ or $S = E_7(q)$. Setting $t = r_8(q)$ in the former case and $t = r_9(q)$ in the latter, we see that $pt \in \omega(H)$ by Lemma 3.2 and $pt \notin \omega(S)$ by [36, Proposition 3.2].

We turn now to symplectic and orthogonal groups. To check whether or not a given integer belongs to the spectrum of S, we use [37].

Let $n \neq 4$ and let S be one of the groups $O_{2n+1}(q)$, with q odd, or one the groups $O^{\pm}_{2n+2}(q)$. Define a number t as follows. If n is even and $n \geqslant 8$, then take integers n_1 and n_2 such that $n = n_1 + n_2$, $n_1 > n_2 \geqslant 3$ and $(n_1, n_2) = 1$, and set $t = r_{n_1}(q)r_{n_2}(q)$. For $n = 6$, set $t = r_3(q)r_6(q)$ if $q \neq 2$ and $t = 63$ otherwise. If n is odd, then $t = r_n(q)$. It is easily seen that $t \in \omega(SL_n(q))$, therefore, $pt \in \omega(H)$ by Lemma 3.2. On the other hand, using [36, Proposition 3.1] when t is a prime and [37] in other cases, we calculate that $pt \notin \omega(S)$.

Let $S = O_9(q), O^{\pm}_{10}(q)$, where q is odd. If $q \equiv \epsilon \pmod 4$, then $p(q^2+1)(q-\epsilon) \notin \omega(S)$. On the other hand, $(q^2+1)(q-\epsilon) \in \omega(SL^{\epsilon}_4(q))$, and so $p(q^2+1)(q-\epsilon) \in \omega(H)$ by Lemma 3.2.

Let $S = O^{\pm}_{10}(q)$, where q is even. Since S includes $Sp_8(q)$, it includes a Frobenius subgroup with cyclic kernel of order q^4+1 and cyclic complement of order 8 by Lemma 2.6. The action of S on V is faithful, and it follows by Lemma 2.4 that H has an element of order 16. It remains to note that $16 \notin \omega(S)$.

Let $S = S_{2n}(q)$, where $n \geqslant 2$ and $q = p^k$ is odd. Take $G = C_n(q)_u$ and identify S with $G/Z(G)$. Denote the root system of G by Φ and enumerate the fundamental roots as in Table 1. Let α_0 be the highest root of Φ. The extended Dynkin diagram of Φ is as follows.



Figure 1: Extended Dynkin diagram of type C_n

Observe that $\alpha_0 = 2(\alpha_1 + \cdots + \alpha_{n-1}) + \alpha_n$, and using [21, Theorem 1.12.1e], we calculate that $h_{\alpha_0}(t) = h_{\alpha_1}(t) \ldots h_{\alpha_n}(t)$. Denote the subgroups generated by the root subgroups associated with the roots $\pm\alpha_2, \ldots, \pm\alpha_n$ and the roots $\pm\alpha_0$ by A and B respectively. Then A and B centralize each other, $A \simeq C_{n-1}(q)_u$ and $B \simeq A_1(q)_u$. Furthermore, a product of the central involutions z_A and z_B of A and B, respectively, is equal to the central involution z of G. By Lemma 3.1, every element of A has a nonzero fixed vector in V. Choose $g \in A$ of order $q^{n-1} - \epsilon$, where $\epsilon = 1$ if $k(n-1)$ is odd and $\epsilon = -1$ otherwise, and set $W = C_V(g)$. The group B centralizes g and so acts on W. Observe that z_B acting trivially because z_A is a power of g and z acts trivially on the whole V. Thus W is a nonzero $B/Z(B)$-module. Embedding $PSL_2(p)$ into $B/Z(B)$ and exploiting unisingularity of $PSL_2(p)$ [31, Theorem 3.12], we can take a element $h \in B$ of order $p+1$ such that $C_W(h) \neq 0$. Since $(q^{n-1} - \epsilon, p+1) = 2$ by choice of ϵ, the image of gh in S has order $(q^{n-1} - \epsilon)(p+1)/2$, and hence $p(q^{n-1} - \epsilon)(p+1)/2 \in \omega(H)$. It is not hard to see that $p(q^{n-1} - \epsilon)(p+1)/2 \notin \omega(S)$.

Let $S = S_{2n}(q)$, where $n \geqslant 4$ and q is even. It is more convenient to consider S as $B_n(q)_u$. Again denote the root system of S by Φ, enumerate the fundamental roots as in Table 1 and denote the highest root of Φ by α_0.



Figure 2: Extended Dynkin diagram of type B_n

The set of long roots in Φ is a subsystem of type D_n, and the subgroup A generated by the corresponding root subgroups is isomorphic to $D_n(q)_u$. Let n be even. Denote the subgroups

generated by the root subgroups associated with the roots $\pm\alpha_1, \pm\alpha_2, \ldots, \pm\alpha_{n-2}$ and the roots $\pm\alpha_n$ by B and C respectively. It is clear that $B \simeq A_{n-2}(q)_u$, $C \simeq A_1(q)_u$ and C centralizes B. By Lemma 3.1, every element of B has a nonzero fixed vector in V. Choose $g \in B$ of order $r_{n-1}(q)$ and set $W = C_V(g)$. The group C acts on W and includes a Frobenius subgroup with cyclic kernel F of order $q+1$ and complement of order 2. If $F \subseteq C_C(W)$, then H has an element of order $2(q+1)r_{n-1}(q)$. Otherwise, Lemma 2.4 yields $4r_{n-1}(q) \in \omega(H)$. None of the numbers $2(q+1)r_{n-1}(q)$ and $4r_{n-1}(q)$ lies in $\omega(S)$. Let n be odd. Then take B and C to be the subgroups generated by the root subgroups associated with $\pm\alpha_1, \pm\alpha_2, \ldots, \pm\alpha_{n-3}$ and $\pm\alpha_{n-1}, \pm\alpha_n$ respectively. Then $B \simeq A_{n-3}(q)_u$, $C \simeq B_2(q)_u$ and again C centralizes B. Choosing an element of order $r_{n-2}(q)$ in B and working with a Frobenius group with cyclic kernel of order q^2+1 and cyclic complement of order 4, which C includes by Lemma 2.6, we conclude that H has either an element of order $2(q^2+1)r_{n-2}(q)$ or an element of order $8r_{n-2}(q)$. And again none of these numbers lies in $\omega(S)$.

The case when $S = S_6(q)$ and q is even requires some additional notation, so we extract this case as a separate lemma. Observe that this lemma is a part of Lemma 3.7 in [38], but the corresponding proof has a gap.

Lemma 4.1. *Let $S = S_6(q)$, where q is even. If V is a nontrivial S-module over a field of characteristic 2 and H is a natural semidirect product of V and S, then $\omega(S) \neq \omega(H)$.*

Proof. Let K be the algebraic closure of the binary field and let $\overline{G} = Sp_6(K)$. Let W be the natural $\overline{G}$-module and f an nondegenerate alternating bilinear form on W preserved by $\overline{G}$. There is a basis $e_1, \ldots, e_6$ of W such that $f(e_i, e_j) = 1$ if $i + j = 7$ and $f(e_i, e_j) = 0$ otherwise. Denote the set of matrices of $\overline{G}$ that are diagonal with respect to this basis by $\overline{D}$. Then $\overline{D}$ is a maximal torus of $\overline{G}$, and ε_1, ε_2, ε_3 defined by the formula $\varepsilon_i(d) = d(e_i)$ for $d \in \overline{D}$ are weights of $\overline{G}$ with respect to this torus. The weights $\lambda_1 = \varepsilon_1$, $\lambda_2 = \varepsilon_1 + \varepsilon_2$, and $\lambda_3 = \varepsilon_1 + \varepsilon_2 + \varepsilon_3$ are the fundamental weights of $\overline{G}$. Given a dominant weight λ, we denote the irreducible $\overline{G}$-module with high weight λ by $L(\lambda)$.

Identify S with $C_{\overline{G}}(\sigma)$, where σ is the Frobenius endomorphism of $\overline{G}$ that raises matrix entries to the q-th power. By [13, Lemma 9], we may assume that the action of S on V is absolutely irreducible. By Steinberg's theorem [39, 13.3], the module V can be obtained as a restriction of some irreducible $\overline{G}$-module $\overline{V}$ to S. Denote the high weight of $\overline{V}$ by λ. We show that $\omega(H) \setminus \omega(S)$ contains an element depending on λ. Note that the even numbers lying in $\omega(S)$ are exactly divisors of 8, $4(q \pm 1)$, and $2(q^2 \pm 1)$.

Suppose that λ is neither $2^j\lambda_1$ nor $2^j\lambda_3$, with $j \geqslant 0$. Then by [40, Theorem 1], for every unipotent element u of $\overline{G}$, the degree of the minimal polynomial of u on $\overline{V}$ is equal to $|u|$. Therefore, for every unipotent element u of S, the degree of the minimal polynomial of u on V is equal to $|u|$ as well. Taking $u \in S$ to be a unipotent element of order 8, we see that the coset Vu has an element of order 16.

Suppose that $\lambda = 2^j\lambda_3$, where $j \geqslant 0$. Let $x \in S$ be an element of order q^2+q+1, and $\zeta \in K$ a primitive (q^2+q+1)-th root of unity. Then x is conjugate to the diagonal matrix y such that $\varepsilon_i(y) = \zeta^{q^{i-1}}$ for $1 \leqslant i \leqslant 3$ in $\overline{G}$. Since $\lambda(y) = \lambda_3(y)^{2^j} = (\zeta^{1+q+q^2})^{2^j} = 1$, it follows that y has a nonzero fixed vector in $\overline{V}$. Then x also has a nonzero fixed vector in $\overline{V}$. This implies that x has a nonzero fixed vector in V, and so the coset Vx contains an element of order $2(q^2+q+1)$.

Lastly, let $\lambda = 2^j\lambda_1$, where $j \geqslant 0$. Assume, for a while, that $j = 0$, and hence V is the natural module of S. Exploiting the standard embedding of $Sp_2(q) \times Sp_4(q)$ into $S_6(q)$, we can find commuting elements $x \in S$ and $y \in S$ of orders 3 and 4 respectively such that $\dim C_V(x) = 4$ and the degree of the minimal polynomial of y on $C_V(x)$ is equal to 4. An easy

computation establishes that the coset Vxy contains an element of order 24. Now let $j > 0$. By Steinberg's tensor product theorem [39], we have $L(\lambda) \simeq L(\lambda_1)^{(\rho^j)}$, where ρ is the Frobenius endomorphism of $\overline{G}$ rasing matrix entries to the second power. Since twisting by ρ does not affect the dimension of the fixed subspace or the degree of the minimal polynomial, the previous argument works, and again H contains an element of order 24.

Thus at least one of the numbers 16, $2(q^2+q+1)$, and 24 lies in $\omega(H) \setminus \omega(S)$, and so the lemma follows.

Finally, let $S = S_4(q)$, where $q > 2$ is even. By Lemma 2.6, there is a Frobenius subgroup with kernel of order q^2+1 and cyclic complement of order 4 in S. It follows that $8 \in \omega(H)\setminus\omega(S)$, and this completes the proof of Proposition 1.1.

Proposition 1.2. *Let S be one of the simple groups ${}^3D_4(q)$, $F_4(q)$, $E_6(q)$, ${}^2E_6(q)$, $E_7(q)$, and $S_4(q)$. If V is a nontrivial S-module over a field of characteristic r, where r is coprime to q, and H is a natural semidirect product of V and S, then $\omega(S) \neq \omega(H)$.*

Proof. Similarly to the proof of Proposition 1.1, we can assume that the action of S on V is faithful. Denote the r-exponent of S by s.

Let S be one of the groups ${}^3D_4(q)$ and $F_4(q)$. By [23,24], the prime graph of S is disconnected with $\pi(q^4-q^2+1)$ being a component. Suppose that $\omega(H) = \omega(S)$. Then the prime graph of H is also disconnected, and by [23, Theorem A], we see that r and 2 lie in the same component of this graph. Since q^4-q^2+1 is odd, it follows that r does not divide q^4-q^2+1 and $r(q^4-q^2+1) \notin \omega(H)$. Exploiting the fact that ${}^3D_4(q)$ is a subgroup of $F_4(q)$ (see, for example, [41]) and applying [42, Proposition 2], we conclude that some element of S of order q^4-q^2+1 has a nonzero fixed vector in V. Thus $r(q^4-q^2+1) \in \omega(H)$, and this is a contradiction.

Let $S = E_6^{\varepsilon}(q)$ and set $d = (3, q-\varepsilon)$. By [43, Theorem 1], the set $\omega_{p'}(S)$ consists of the divisors of the numbers

$$\frac{q^6+\varepsilon q^3+1}{d}, \frac{(q^4-q^2+1)(q^2+\varepsilon q+1)}{d}, \frac{(q^5-\varepsilon)(q+\varepsilon)}{d}, q^5-\varepsilon,$$
$$\frac{(q^4+1)(q^2-1)}{d}, \frac{q^6-1}{d}, (q^3-\varepsilon)(q+\varepsilon), \frac{(q^4-1)(q^2-\varepsilon q+1)}{d}, q^4-1. \quad (1)$$

Let $\varepsilon = +$. The universal group $E_6(q)_u$ contains a subsystem subgroup A isomorphic $A_5(q)_u$. By Lemma 2.5, the group A contains Frobenius subgroups of orders $q^5(q^5-1)_{(6,q-1)'}$ and $q^4(q^4-1)$. Clearly, these subgroups meet $Z(E_6(q)_u)$ trivially, and so they can be embedded into S. Then by Lemma 2.4, there are elements of orders $r(q^5-1)_{(6,q-1)'}$ and $r(q^4-1)$ in H. Suppose that both of these numbers lie in $\omega(S)$. Examining the numbers in (1) and using Zsigmondy's theorem, we calculate that r divides $(q^5-1)(q+1)$ and q^2-q+1. Since $((q^5-1)(q+1), q^2-q+1) = (q+1, 3)$, it follows that r is equal to 3 and it divides $q+1$. Then $s = (q^3+1)_3$. A maximal parabolic subgroup of S with Levi factor of type $D_5(q)$ has an abelian unipotent radical (see, for example [44, Theorem 1]) and $s \in \omega(D_5(q))$, therefore, by Lemma 2.7, there is an element of order $3s$ in H. Thus $\omega(H) \neq \omega(S)$.

Let $\varepsilon = -$. Considering a subsystem subgroup $A_2(q^2)_u$ in ${}^2E_6(q)_u$ and applying Lemma 2.5, we conclude that S has a Frobenius subgroup of order $q^4(q^4-1)_{(3,q^2-1)'}$. By Lemma 2.4 again, we have $r(q^4-1)_{(3,q^2-1)'} \in \omega(H)$. Suppose that $r(q^4-1)_{(3,q^2-1)'} \in \omega(S)$. Working with the 2-parts of the numbers in (1), we deduce that r divides $(3, q^2-1)(q^2+q+1)$, and in particular, r is odd. Assume that r divides q^2+q+1. Then $s = (q^3-1)_r$. Consider a subsystem subgroup of ${}^2E_6(q)_u$ isomorphic to ${}^2A_5(q)_u$. Thus subgroup includes $Z({}^2E_6(q)_u)$ (see Table 1), and hence

its image in S is an extension of the group of order $(2, q-1)$ by ${}^2A_5(q)$. By [14, Lemma 5(4)], the group ${}^2A_5(q)$ contains a Frobenius subgroup with kernel of order q^3 and cyclic complement of order $(q^3-1)_{(2,q-1)'}$. It follows that S also contains such a Frobenius subgroup, and so $r(q^3-1)_{2'} \in \omega(H)$. Thus $rs \in \omega(H) \setminus \omega(S)$. Now assume that r does not divide q^2+q+1. Then $r=3$ and it divides q^2-1. If $q-1$ is divisible by 3, then so is q^2+q+1, contrary to our assumption. So 3 divides $q+1$ and $s=(q^3+1)_3$. A maximal parabolic subgroup of S with Levi factor of type ${}^2D_4(q)$ contains an element of order s, therefore, if q is odd, then we can apply Lemma 2.7 to deduce that $rs \in \omega(H) \setminus \omega(S)$. If q is even, then S contains a subgroup isomorphic to ${}^2D_4(q)$. A maximal parabolic subgroup of ${}^2D_4(q)$ with Levi factor of type ${}^2A_3(q)$ has an abelian unipotent radical (see, for example, [34, Table 8.52]) and contains an element of order s. Applying Lemma 2.7, we see that $rs \in \omega(H) \setminus \omega(S)$ in this case too.

Let $S = E_7(q)$. The structure of maximal tori of $E_7(q)_u$ described in [45] implies that $\omega_{p'}(E_7(q)_u)$ consists of the divisors of the numbers

$$(q^6+\epsilon q^3+1)(q-\epsilon), q^7-\epsilon, (q^4-q^2+1)(q^3-\epsilon), (q^5-\epsilon)(q^2+\epsilon q+1),$$
$$(q^5-\epsilon)(q+\epsilon), \frac{(q^4+1)(q^2+1)(q-\epsilon)}{(2,q-1)}, (q^4+1)(q^2-1), (q^4-1)(q^2+\epsilon q+1), q^6-1, \quad (2)$$

where ϵ runs over $\{+1,-1\}$. Since $E_7(q)$ contains a quotient of $SL_8(q)$ by a central 2-subgroup, the group S contains Frobenius subgroups of orders $q^7(q^7-1)_{(2,q-1)'}$ and $q^4(q^4-1)$. It follows that $r(q^7-1)_{(2,q-1)'}$ and $r(q^4-1)$ lie in $\omega(H)$. On the other hand, examining the numbers in (2), we see that $r(q^4-1) \notin \omega(S)$ if $r=2$ and $r(q^7-1)_{(2,q-1)'} \notin \omega(S)$ if r is odd.

Lastly, let $S = S_4(q)$, where $q > 2$. By [46, Lemma 2.8], there is a Frobenius subgroup with kernel of order q^2 and cyclic complement of order $(q^2-1)/(2,q-1)$ in S. By Lemma 2.4 yet again, we have $r(q^2-1)/(2,q-1) \in \omega(H)$. By [37], the number $(q^2-1)/(2,q-1)$ is maximal under divisibility in $\omega(S)$, and the proof of Proposition 1.2 is complete.

Now we are ready to prove the theorem. Let S and H be as in the hypothesis and let S be defined over a field of characteristic p. By Lemma 2.3, we can assume that H is a semidirect product of a nonzero finite S-module over a field of characteristic r by S. If $r=p$, then $\omega(H) \neq \omega(S)$ by Proposition 1.1. Let $r \neq p$. If S is an orthogonal or symplectic group other than $S_4(q)$, then $\omega(H) \neq \omega(S)$ by [14, Theorem 1]. Otherwise, $\omega(H) \neq \omega(S)$ by Proposition 1.2, and the theorem follows.

5. Covers of $O_{2n+1}(q)$ and the spectrum of $S_{2n}(q)$

A simple nonabelian group L is said to be *quasirecognizable by spectrum* if every finite group G with $\omega(G) = \omega(L)$ has exactly one nonabelian composition factor and this factor is isomorphic to L. The following result was obtained in studying quasirecognizability of simple symplectic groups: If $L = S_{2n}(q)$, where $n \geqslant 4$, and G is a finite group such that $\omega(G) = \omega(L)$, then G has exactly one nonabelian composition factor and if this factor is a group of Lie type in the same characteristic as L, then it is isomorphic either to L or to one of the groups $O_{2n+1}(q)$ and $O^-_{2n}(q)$ [20, Theorem 3]. As we mentioned in Introduction, the case when the factor is isomorphic to $O_{2n+1}(q)$ requires a separate treatment because the spectra of $S_{2n}(q)$ and $O_{2n+1}(q)$ are very close.

Proposition 1.3. *Let $L = S_{2n}(q)$ and $S = O_{2n+1}(q)$, where $n \geqslant 3$ and q is odd. If H is a proper cover of S, then $\omega(H) \not\subseteq \omega(L)$. In particular, if G is a finite group such that $\omega(G) = \omega(L)$ and G has a composition factor isomorphic to S, then $S \leqslant G \leqslant \mathrm{Aut}(S)$.*

Proof. Let q be a power of a prime p. By Lemma 2.3, we can assume that $H = V \leftthreetimes S$, where V is a nonzero S-module over a field of a positive characteristic r. Suppose that $r \neq p$ and $(S, r) \neq (O_7(3), 5)$. Then by the proof of Proposition 4 in [14], the group H contains an element of order rt such that p is coprime to t and $rt \notin \omega(S)$. Since the isomorphism types of maximal tori in S are the same as in L [47], we have $\omega_{p'}(S) = \omega_{p'}(L)$, and so $rt \notin \omega(L)$. If $S = O_7(3)$ and $r = 5$, then the 5-Brauer character table of S [48] implies that every element of S of order 7 has a nonzero fixed vector in V, and hence $35 \in \omega(H) \setminus \omega(L)$. Now let $r = p$. If $n \neq 4$, then we showed in the proof of Proposition 1.1 that H contains an element of order pt, where t is odd and $pt \notin \omega(S)$. One can show (by using, for example, [37]) that the subsets of odd numbers are the same in $\omega(S)$ and $\omega(L)$, and thus $pt \notin \omega(L)$. If $n = 4$ and $q \equiv \epsilon \pmod 4$, then we proved that $p(q^2+1)(q-\epsilon) \in \omega(H)$. By [37], we see that $p(q^2+1)(q-\epsilon) \notin \omega(L)$.

Now suppose that G is a finite group such that $\omega(G) = \omega(L)$. By [36, Corollary 7.2], the group G has exactly one nonabelian composition factor and this factor is isomorphic to S. Denoting the soluble radical of G by K, we see that $S \leqslant G/K \leqslant \mathrm{Aut}(S)$. If $K \neq 1$, then G contains a proper cover of S and so $\omega(G) \not\subseteq \omega(L)$, which contradicts the hypothesis. Thus $K = 1$, and the proof is complete.

References

[1] V. D. Mazurov, Recognition of finite groups by a set of orders of their elements, *Algebra and Logic* **37** (1998) 371–379.

[2] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker and R. A. Wilson, *Atlas of finite groups*, (Clarendon Press, Oxford, 1985).

[3] V. D. Mazurov and W.J. Shi, A note to the characterization of sporadic simple groups, *Algebra Colloq.* **5** (1998) 285–288.

[4] A. V. Zavarnitsine and V. D. Mazurov, Element orders in coverings of symmetric and alternating groups, *Algebra and Logic* **38** (1999) 159–170.

[5] W.J. Shi, A characterization of Suzuki's simple groups, *Proc. Amer. Math. Soc.* **114** (1992) 589–591.

[6] R. Brandl and W.J. Shi, A characterization of finite simple groups with abelian Sylow 2-subgroups, *Ricerche Mat.* **42** (1993) 193–198.

[7] H.W. Deng and W.J. Shi, The characterization of Ree groups ${}^2F_4(q)$ by their element orders, *J. Algebra* **217** (1999) 180–187.

[8] A. V. Vasil'ev and A. M. Staroletov, Recognizability of $G_2(q)$ by spectrum, *Algebra Logic* **52** (2013) 1–14.

[9] A. S. Kondrat'ev, Recognizability by spectrum of $E_8(q)$, *Trudy Inst. Mat. Mekh. UrO RAN* **16** (2010) 146–149.

[10] R. Brandl and W.J. Shi, The characterization of $PSL(2, q)$ by its element orders, *J. Algebra* **163** (1994) 109–114.

[11] A. V. Zavarnitsine, Recognition of the simple groups $L_3(q)$ by element orders, *J. Group Theory* **7** (2004) 81–97.

[12] A. V. Zavarnitsine, The weights of irreducible $SL_3(q)$-modules in the defining characteristic, *Siberian Math. J.* **45** (2004) 261–268.

[13] A. V. Zavarnitsine, Properties of element orders in covers for $L_n(q)$ and $U_n(q)$, *Siberian Math. J.* **49** (2008) 246–256.

[14] M. A. Grechkoseeva, On element orders in covers of finite simple classical groups, *J. Algebra* **339** (2011) 304–319.

[15] V. D. Mazurov and G.Y. Chen, Recognizability of the finite simple groups $L_4(2^m)$ and $U_4(2^m)$ by the spectrum, *Algebra Logic* **47** (2008) 49–55.

[16] A. V. Zavarnitsine, Recognition of the simple groups $U_3(q)$ by element orders, *Algebra Logic* **45** (2006) 106–116.

[17] A. V. Zavarnitsine, Exceptional action of the simple groups $L_4(q)$ in the defining characteristic, *Sib. Élektron. Mat. Izv.* **5** (2008) 65–74.

[18] V. D. Mazurov, Unrecognizability by spectrum for a finite simple group ${}^3D_4(2)$, *Algebra Logic* **52** (2013) 400–403.

[19] V. D. Mazurov and E.I. Khukhro, eds., *The Kourovka notebook. Unsolved problems in group theory*, (Institute of Mathematics, Russian Academy of Sciences, Siberian Div., Novosibirsk, 2010), 17 edition.

[20] A. V. Vasil'ev, M. A. Grechkoseeva and V. D. Mazurov, On finite groups isospectral to simple symplectic and orthogonal groups, *Siberian Math. J.* **50** (2009) 965–981.

[21] D. Gorenstein, R. Lyons and R. Solomon, *The classification of the finite simple groups. Number 3*, *Mathematical Surveys and Monographs*, volume 40.3, (American Mathematical Society, Providence, RI, 1998).

[22] K. Zsigmondy, Zur Theorie der Potenzreste, *Monatsh. Math. Phys.* **3** (1892) 265–284.

[23] J. S. Williams, Prime graph components of finite groups, *J. Algebra* **69** (1981) 487–513.

[24] A. S. Kondrat'ev, On prime graph components of finite simple groups, *Math. USSR-Sb.* **67** (1990) 235–247.

[25] A. V. Vasil'ev and E. P. Vdovin, Cocliques of maximal size in the prime graph of a finite simple group, *Algebra Logic* **50** (2011) 291–322.

[26] V. D. Mazurov, Characterizations of finite groups by sets of orders of their elements, *Algebra and Logic* **36** (1997) 23–32.

[27] A. V. Vasil'ev and M. A. Grechkoseeva, On recognition by spectrum of finite simple linear groups over fields of characteristic 2, *Siberian Math. J.* **46** (2005) 593–600.

[28] A. V. Vasil'ev and M. A. Grechkoseeva, On recognition of the finite simple orthogonal groups of dimension 2^m, 2^{m+1}, and 2^{m+2} over a field of characteristic 2, *Siberian Math. J.* **45** (2004) 420–432.

[29] P. Hall and G. Higman, On the p-length of p-soluble groups and reduction theorem for Burnside's problem, *Proc. London Math. Soc.* **6** (1956) 1–42.

[30] I. D. Suprunenko and A. E. Zalesski, Fixed vectors for elements in modules for algebraic groups, *Int. J. Algebra Comput.* **17** (2007) 1249–1261.

[31] L. Babai and A. Shalev, Recognizing simplicity of black-box groups and the frequency of p-singular elements in affine groups, *Groups and computation, III (Columbus, OH, 1999)*, (de Gruyter, Berlin, 2001), *Ohio State Univ. Math. Res. Inst. Publ.*, volume 8, 39–62.

[32] D. I. Deriziotis, *Conjugacy classes of centralizers of semisimple elements in finite groups of Lie type*, *Vorlesungen Fachbereich Math. Univ. Essen*, volume 11, (Universität Essen Fachbereich Mathematik, Essen1984).

[33] R.M. Guralnick and P.H. Tiep, Finite simple unisingular groups of Lie type, *J. Group Theory* **6** (2003) 271–310.

[34] J. Bray, D. Holt and C. Roney-Dougal, *The maximal subgroups of the low-dimensional finite classical groups*, *London Mathematical Society Lecture Note Series*, volume 407, (Cambridge University Press, Cambridge, 2013).

[35] D. I. Deriziotis and G. O. Michler, Character table and blocks of finite simple triality groups ${}^3D_4(q)$, *Trans. Amer. Math. Soc.* **303** (1987) 39–70.

[36] A. V. Vasil'ev and E.P. Vdovin, An adjacency criterion for the prime graph of a finite simple group, *Algebra Logic* **44** (2005) 381–406.

[37] A. A. Buturlakin, Spectra of finite symplectic and orthogonal groups, *Siberian Adv. Math.* **21** (2011) 176–210.

[38] A. M. Staroletov, On recognition by spectrum of the simple groups $B_3(q)$, $C_3(q)$ and $D_4(q)$, *Siberian Math. J.* **53** (2012) 532–538.

[39] R. Steinberg, *Endomorphisms of linear algebraic groups*, number 80 in Memoirs of the American Mathematical Society, (American Mathematical Society, Providence, RI, 1968).

[40] I.D. Suprunenko, The minimal polynomials of unipotent elements in irreducible representations of the symplectic group over a field of characteristic two, *Dokl. Nats. Akad. Nauk Belarusi* **48** (2004) 28–31.

[41] E. Stensholt, Certain emdeddings among finite groups of Lie type, *J. Algebra* **53** (1978) 136–187.

[42] A. V. Zavarnitsine, Finite groups with a five-component prime graph, *Siberian Math. J.* **54** (2013) 40–46.

[43] A. A. Buturlakin, Spectra of finite simple groups $E_6(q)$ and ${}^2E_6(q)$, *Algebra Logic* **52** (2013) 188–202.

[44] A. V. Vasilyev, Minimal permutation representations of finite simple exceptional groups of types E_6, E_7 and E_8, *Algebra and Logic* **36** (1997) 302–310.

[45] D. I. Deriziotis and A. P. Fakiolas, The maximal tori in the finite Chevalley groups of type E_6, E_7 and E_8, *Commun. Algebra* **19** (1991) 889–903.

[46] H. He and W. J. Shi, Recognition of some finite simple groups of type $D_n(q)$ by spectrum, *Int. J. Algebra Comput.* **19** (2009) 681–698.

[47] R. W. Carter, Centralizers of semisimple elements in the finite classical groups, *Proc. Lond. Math. Soc. (3)* **42** (1981) 1–41.

[48] C. Jansen, K. Lux, R. Parker and R. Wilson, *An atlas of Brauer characters*, *London Mathematical Society Monographs. New Series*, volume 11, (Clarendon Press, 1995).